

# Is Large-Scale Low-Frequency Variability Satisfying Linear Barotropic Vorticity Dynamics?

Lee-Lueng Fu and Roger Davidson

Jet Propulsion Laboratory, California Institute of Technology  
Pasadena, CA 91109 USA

What is the dynamics governing the sea level variabilities at spatial scales larger than the **mesoscale** and periods longer than a few weeks? This question is investigated using the **TOPEX/POSEIDON** altimetry **data**. For periods shorter than a decade, **it** is believed that the large-scale variabilities in the open ocean should be **barotropic** and can be described by the **barotropic vorticity** equation:

$$\frac{\partial}{\partial t} \nabla^2 \eta + \beta \frac{\partial \eta}{\partial x} - \frac{f}{H} \left( \frac{\partial \eta}{\partial x} \frac{\partial H}{\partial y} - \frac{\partial \eta}{\partial y} \frac{\partial H}{\partial x} \right) = \frac{f}{\rho g H} (\nabla \times \tau)_z$$

where  $\eta$  is the sea level height,  $H$  the ocean depth,  $f$  the Coriolis parameter,  $\beta = df/dy$ ,  $\rho$  the water density,  $g$  the gravity,  $\tau$  the wind stress. To avoid the error-sensitive **Laplacian** operator, the above equation was integrated over a large area. Preliminary analysis was performed in the northeastern Pacific, where the eddy energy is relatively low and the **bathymetry** is relatively -th. Good correlation between the two **sides of** the equation was obtained at periods **longer** than 60 **days**. The **largest error in the** data is suspected **to** be the ocean tides. **Empirical** correction for **the** ocean tides will be performed **for** further analysis. Preliminary results **of a global** calculation **will** be presented.